



DRYMODES BRUNNEOPAGA: *Gould*

## DRYMODES BRUNNEOPYGIA, *Gould.*

### Scrub Robin.

*Drymodes brunneopygia*, Gould in Proc. of Zool. Soc., Part VIII. p. 170.

---

I DISCOVERED this singular bird in the great Murray Scrub in South Australia, where it was tolerably abundant; I have never seen it from any other part of the country, and it is doubtless confined to such portions of Australia as are clothed with a similar character of vegetation. It is a quiet and inactive species, resorting much to the ground, over which and among the underwood and low stunted bushes it passes with great ease; it appeared rarely to take wing, but to depend for security upon its dexterity in hopping away under the dense underwood of the most scrubby parts; I have, however, occasionally observed it to mount to the most elevated part of a low bush, and there pour forth a sharp monotonous whistling note, not very unlike that of some of the *Pachycephalinæ*; indeed it was its note that first attracted my attention and led to its discovery. When on the ground, and sometimes when perched on a twig, it elevates its tail considerably, but not to the extent of the *Mahuri*.

This new form evidently belongs to the *Saxicolinæ*, and has many habits in common with the members of the genus *Petroica*.

The sexes are alike in colouring, but the female is much smaller than her mate; the young, as will be seen in the accompanying Plate, resembles the immature *Petroicas* in the character of its plumage.

Head and all the upper surface brown, passing into rufous brown on the upper tail-coverts; wings dark brown, the coverts and primaries edged with dull white; primaries and secondaries crossed near the base on their inner webs with pure white; tail rich brown, all but the two middle feathers tipped with white; under surface greyish brown, passing into buff on the under tail-coverts; irides, bill and feet blackish brown.

The Plate represents an adult male and a young bird of the natural size.